

NOTES AND FIELD REPORTS

Chelonian Conservation and Biology, 2000, 3(4):658–660
© 2000 by Chelonian Research Foundation

Predation of Neonate Blanding's Turtles (*Emydoidea blandingii*) by Short-Tailed Shrews (*Blarina brevicauda*)

K. LORRAINE STANDING¹, THOMAS B. HERMAN^{1,3},
AND IAN P. MORRISON²

¹Centre for Wildlife and Conservation Biology, Acadia
University, Wolfville, Nova Scotia, B0P 1X0 Canada;

²Parks Canada, Kejimikujik National Park, Maitland Bridge,
Annapolis County, Nova Scotia, B0T 1B0 Canada;

³Corresponding Author for Reprint Requests
[Fax: 902-585-1059; E-mail: tom.herman@acadiu.ca]

Blanding's turtle (*Emydoidea blandingii*) is a North American freshwater species with a range centered south of the Great Lakes (Herman et al., 1995). Isolated populations occur outside the main range in Wisconsin (Cochran and Lyons, 1986; Ross, 1989), Massachusetts (Butler and Graham, 1995), New York (Breisch, 1997; Kiviat, 1997), Minnesota (Sajwaj et al., 1998), New Hampshire (Taylor, 1997), Maine (McCullough, 1997), and Nova Scotia (Herman et al., 1995). Populations in Nova Scotia have been designated nationally Threatened (Herman et al., 1995) and provincially Endangered (Nova Scotia Government, 2000); recent research is being applied to the development of conservation strategies appropriate for this population (Herman et al., 1999).

Like many turtles, Blanding's turtles are long-lived (Brecke and Moriarty, 1989; Congdon and van Loben Sels, 1993; Herman et al., 1995), reach sexual maturity late (Petokas, 1986; Congdon et al., 1993), and populations are vulnerable to increases in both juvenile and adult mortality (Iverson, 1991; Congdon et al., 1993). In Nova Scotia, concern over the scarcity of sexually immature Blanding's turtles and the low recruitment into the breeding population contributed to the status designation for this population (Herman et al., 1995). A nest protection program was implemented in Kejimikujik National Park, Nova Scotia, in order to bolster recruitment by mitigating nest destruction by predators, and to learn more about other factors influencing clutch failure in this population (Herman et al., 1999; Standing et al., 2000). To date, nest protection is the most important aspect of the headstart program for this population, although an experimental evaluation of the feasibility of captive rearing and release of headstarted hatchlings is currently underway. Since the objective of hatchling headstart programs is to raise neonates to a size sufficiently large to reduce their vulnerability to predation, the identification of local predator species is of interest. Such information will help managers determine the minimum size requirements necessary for an effective headstart program. We report observations on the predation of neonate Blanding's turtles (*E. blandingii*) by short-

tailed shrews (*Blarina brevicauda*) in Kejimikujik National Park, Nova Scotia, Canada.

Methods and Results. — Data were collected in 1994 and 1995 during a study of the early postemergence behavior of neonates in this population (Standing et al., 1997). Detailed descriptions of the study site are available elsewhere (Power, 1989; Power et al., 1994; Herman et al., 1995; McNeil, 1996; McMaster, 1996; Standing et al., 1997).

During the nesting season (June and July), 23 freshly laid Blanding's turtle nests were protected against predation. A solid wood frame (50 cm² x 10 cm high) screened with one-inch (2.5 cm) hardware cloth was centered over each nest. The box frame was placed in a shallow trench dug around the nest, such that the screen was about 10 cm above the nest surface. The protective box was secured in place by large rocks. These screened boxes effectively guarded against predation of eggs and served as pens for emergent hatchlings.

Hatchling emergence occurred throughout September and October. Within clutches, hatchling emergence was mostly asynchronous (i.e., occurring over several days). Hatchlings emerged during the day and were diurnally active. Upon emergence, numerous hatchlings were measured, weighed, powdered with tracking-pigment, and their trails tracked from 1 to 11 days after release (Standing et al., 1997). Most hatchlings were released on the day of emergence, although some hatchlings spent one night after emergence in the enclosures.

After release, hatchlings used terrestrial and aquatic forms (sleeping and resting sites) both during the day and overnight. While in terrestrial forms neonates usually were well concealed beneath vegetation and roots, or between beach cobble, although sometimes hatchlings remained completely exposed on the beach overnight (McNeil, 1996; Standing et al., 1997).

In 1994, five hatchlings from two nests were found dead at the end of their trails. Four of these were nestmates that were found near the nest of origin shortly after their release. These hatchlings were severely bitten by ants and they presumably died from these bites since their carcasses were intact when found. A fifth hatchling, from another nest, was released at ca. 1300 hrs on 25 September. Shortly after having been located alive at the end of its trail at ca. 2200 hrs, it was seized by a predator. Although its carcass was not found, we heard the predator emerge from nearby shrubs and concluded that it was a medium-sized mammal, possibly a raccoon (*Procyon lotor*). Four additional, unpowdered hatchlings from three nests were depredated while still in their screened enclosures and two others survived with signs of attempted predation (i.e., superficial bites taken from their shells).

Between 19 and 25 September 1995, eviscerated carcasses of four powdered hatchling were found at, or near, the end of their marked trails. These hatchlings were nestmates and were within 20 m of their nest of origin when killed; predation occurred up to 6 days after release. On 19, 20, and 23 September, three additional unpowdered hatchlings from the same clutch were depredated inside the protective nest

screen. On the night of 19 September, shortly after finding one eviscerated hatchling within the nest enclosure, we interrupted a predator while it ate one of the released, powdered hatchlings (at ca. 2040 hrs). While eating this hatchling the predator became covered in tracking powder and we followed its trail for a short distance. Footprints were discernible and were identified as those of the short-tailed shrew (*Blarina brevicauda*).

The most prevalent form of mutilation of these hatchlings was evisceration. Typically, a section of the plastron was eaten, beginning at the axillary scutes, proceeding medially to the midline suture of the abdominal scute, and posteriorly to the inguinal region. Otherwise, a disk centered on the umbilical region was removed. Hatchlings were also eviscerated through the carapace. Either a circular area centered on the suture between the second and third vertebral scutes and extending laterally to the middle of the adjoining costal scutes was removed, or a crescent extending medially from the axillary and inguinal regions was removed. One hatchling that had been eviscerated through the plastron also had the foot and skin eaten from its left hind leg. Two hatchlings were decapitated. Those hatchlings that survived attack had bites taken from the carapace and marginal scutes 5, 6, and 7.

Discussion. — Turtles, including Blanding's turtle, are characterized by a Type III survivorship curve in which the rate of mortality is inversely related to age (Iverson, 1991). Presumably, this results from intense predation on early life stages, particularly eggs (Iverson, 1991). Numerous authors have attributed high clutch failure in *E. blandingii* and other freshwater turtles (*Chelydra serpentina*, *Chrysemys picta*, *Clemmys insculpta*, *Kinosternon flavescens*) to predation by raccoons (*P. lotor*), skunks (*Mephitis mephitis*), foxes (*Vulpes fulva* and *Urocyon cinereoargenteus*), badgers (*Taxidea taxus*), hognose snakes (*Heterodon nasicus*), rodents, and small, unidentified mammals (Congdon et al., 1983; Petokas, 1986; Power, 1989; Iverson, 1990; Ross and Anderson, 1990; Brooks et al., 1991; Herman et al., 1995). While nest screening programs such as the one implemented in Nova Scotia improve clutch survival, these programs are, in themselves, inadequate conservation measures (Heppell et al., 1996). Nest screening is only one element of a headstart program; threatened turtle populations would also benefit by the headstarting of hatchlings and from the protection of sexually immature turtles in the wild (Iverson, 1991; Heppell et al., 1996). In addition to bolstering recruitment, nest screening facilitates the collection of accurate data of hatching success (Standing, 1997). In addition, headstarted hatchlings may help researchers locate and describe important habitat used by immature turtles and actually lead to juvenile turtles in the wild.

It is widely accepted that, like eggs, small turtles are most vulnerable to predation pressure. Since this is the underlying premise of headstart programs (Heppell et al., 1996), the design of effective management strategies requires a thorough knowledge of causes of mortality in small size classes. Confirmed predators of hatchling freshwater turtles of various species include gulls (*Larus* spp.), crows

(*Corvus* spp.), raccoons (*P. lotor*), bullfrogs (*Rana catesbeiana*) (Lefevre and Brooks, 1995), water snakes (*Nerodia* sp.) (Janzen et al., 1992), coyotes (*Canis latrans*) (Minckley, 1966), American Kestrel (*Falco sparverius*) (B. Butler, *pers. comm.*), and larger turtles (Sloan et al., 1996); putative predators include dogs, cats, toads, bears (Ehrenfeld, 1979), unknown shrew species (T. Graham, *pers. comm.*), squirrels (J. McNeil, *pers. comm.*), mink, otter, wading birds, and large, predatory fish (Congdon et al., 1992). Our data indicate that the short-tailed shrew (*B. brevicauda*) should be added to the growing list of predators of small freshwater turtles.

Short-tailed shrews (*B. brevicauda*) are common throughout Nova Scotia. They are opportunistic predators that primarily feed on invertebrates, although they have been known to catch and eat small lizards and mammals (Churchfield, 1990). It is reasonable to suspect them of being able to eat small, hatchling Blanding's turtles, especially since the shells of neonates are not heavily calcified or ossified. Though the foraging of *B. brevicauda* tends to be undirected, shrews will return to a concentration of prey until the supply is exhausted (Churchfield, 1980) and this might explain the apparent exploitation of individual nests in our study. In addition to the style of mutilation and the observed identifiable footprints, other evidence suggestive of predation by *B. brevicauda* is that short-tailed shrews are small enough to fit through one-inch hardware cloth; that is, without disturbing or digging beneath the screened predator enclosures, shrews could have accomplished the observed predation of hatchlings within the enclosures.

As well as providing necessary information for the development of effective headstart programs, identifying species that prey on hatchling freshwater turtles provides insight into the behavior (Janzen et al., 1992; Butler and Graham, 1995; Lefevre and Brooks, 1995) and habitat selection (Congdon et al., 1992; Pappas and Brecke, 1992) of neonate freshwater turtles, and their antipredator mechanisms (Britson and Gutzke, 1993). Combined, such information will help in the development of comprehensive management plans for the protection of young, vulnerable age classes in the wild.

Acknowledgments. — We gratefully acknowledge funding from World Wildlife Fund Canada - Endangered Species Recovery Fund, The Linnaeus Fund and Chelonian Research Foundation, Acadia University Faculty Association Article 25.55 Fund, Nova Scotia Liquor Commission / Nova Scotia Department of the Environment Endangered Species T-shirt Program, and Parks Canada. We also thank Parks Canada for logistical support, and the staff of Kejimikujik National Park and students of Acadia University for their assistance in gathering field data. We also thank J. McNeil, T. Graham, and B. Butler for their communications, and S. Bondrup-Nielsen, S. Boates, F. Scott, T. Graham, B. Butler, and A. Rhodin for their suggestions and criticisms of the manuscript. This study was completed by KLS in partial fulfilment of the requirements for the Degree of Master of Science.

LITERATURE CITED

- BRITSON, C.A. AND GUTZKE, W.H.N. 1993. Antipredator mechanisms of hatchling freshwater turtles. *Copeia* 1993:435-440.
- BRECKE, B. AND MORIARTY, J.J. 1989. Life history notes. *Emydoidea blandingii*. Longevity. *Herp. Rev.* 20:53.
- BREISCH, A. 1997. The status and management of turtles in New York. In: Tying, T.F. (Ed.). Status and Conservation of Turtles of the Northeastern United States. Lanesboro, MN: Serpent's Tale, pp. 11-14.
- BROOKS, R.J., SHILTON, C.M., BROWN, G.P., AND QUINN, N.W.S. 1991. Body size, age distribution, and reproduction in a northern population of wood turtles (*Clemmys insculpta*). *Can. J. Zool.* 70:462-469.
- BUTLER, B.O. AND GRAHAM, T.E. 1995. Early post-emergent behavior and habitat selection in hatchling Blanding's turtles, *Emydoidea blandingii*, in Massachusetts. *Chelonian Conservation and Biology* 1:187-196.
- CHURCHFIELD, S. 1980. Subterranean foraging and burrowing activity of the common shrew. *Acta Theriol.* 25:451-459.
- CHURCHFIELD, S. 1990. The Natural History of Shrews. New York: Comstock Publishing Associates, 178 pp.
- COCHRAN, P.A. AND LYONS, J.D. 1986. New distributional records for Wisconsin amphibians and reptiles. *Trans. Wis. Acad. Sci. Arts Lett.* 74:138-141.
- CONGDON, J.D. AND VAN LOBEN SELS, R.C. 1993. Relationships of reproductive traits and body size with attainment of sexual maturity and age in Blanding's turtles (*Emydoidea blandingii*). *J. Evol. Biol.* 6:547-557.
- CONGDON, J.D., TINKLE, D.W., BREITENBACH, G.L., AND VAN LOBEN SELS, R.C. 1983. Nesting ecology and hatching success in the turtle *Emydoidea blandingii*. *Herpetologica* 39(4):417-429.
- CONGDON, J.D., GOTTE, S.W., AND MCDIARMID, R.W. 1992. Ontogenetic changes in habitat use by juvenile turtles, *Chelydra serpentina* and *Chrysemys picta*. *Can. Field-Nat.* 106:241-248.
- CONGDON, J.D., DUNHAM, A.E., AND VAN LOBEN SELS, R.C. 1993. Delayed sexual maturity and demographics of Blanding's turtles (*Emydoidea blandingii*): implications for conservation and management of long-lived organisms. *Conservation Biol.* 7:826-833.
- EHRENFELD, D.W. 1979. Behavior associated with nesting. In: Harless, M. and Morlock, H. (Eds.). *Turtles: Perspectives and Research*. New York: John Wiley and Sons, pp. 417-434.
- HEPPELL, S.S., CROWDER, L.B., AND CROUSE, D.T. 1996. Models to evaluate headstarting as a management tool for long-lived turtles. *Ecological Applications* 6:556-565.
- HERMAN, T.B., POWER, T.D., AND EATON, B.R. 1995. Status of Blanding's turtle, *Emydoidea blandingii*, in Nova Scotia, Canada. *Can. Field-Nat.* 109:182-191.
- HERMAN, T.B., BLEAKNEY, J.S., BOATES, J.S., DRYSDALE, C., GILHEN, J., MORRISON, I.P., POWER, T.D., STANDING, K.L., AND ELDERKIN, M. 1999. National recovery plan for Blanding's turtle (*Emydoidea blandingii*) Nova Scotia population. Ottawa: Recovery of Nationally Endangered Wildlife Committee, Report No. 18, 39 pp.
- IVERSON, J.B. 1990. Nesting and parental care in the mud turtle, *Kinosternon flavescens*. *Can. J. Zool.* 68:230-233.
- IVERSON, J.B. 1991. Patterns of survivorship in turtles (order Testudines). *Can. J. Zool.* 69:385-391.
- JANZEN, F.J., PAUKSTIS, G.L., AND BRODIE, E.D., III. 1992. Observations on basking of hatchling turtles in the wild. *J. Herpetol.* 26:217-219.
- KIVIAT, E. 1997. Blanding's turtle habitat requirements and implications for conservation in Dutchess County, New York. In: Van Abbema, J. (Ed.). *Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles – An International Conference*. N.Y. Turtle and Tortoise Society, pp. 377-382.
- LEFEVRE, K. AND BROOKS, R.J. 1995. Effects of sex and body size on basking behavior in a northern population of the painted turtle, *Chrysemys picta*. *Herpetologica* 51(2):217-224.
- MCCOLLOUGH, M. 1997. Status and conservation of turtles in Maine. In: Tying, T.F. (Ed.). Status and Conservation of Turtles of the Northeastern United States. Lanesboro, MN: Serpent's Tale, pp. 7-11.
- MCMASTER, N.L. 1996. Age structure, distribution, habitat selection and movement patterns of juvenile Blanding's turtles (*Emydoidea blandingii*) in Kejimikujik National Park, Nova Scotia. B.S. Honours Thesis, Acadia University, Wolfville.
- MCCNEIL, J. 1996. Post-emergent movement of hatchling Blanding's turtles (*Emydoidea blandingii* (Holbrook)) in Kejimikujik National Park, Nova Scotia. B.S. Honours Thesis, Acadia University, Wolfville.
- MINCKLEY, W.L. 1966. Coyote predation on aquatic turtles. *J. Mammal.* 47:137.
- NOVA SCOTIA GOVERNMENT. 2000. N.S. Reg. 109/2000. Endangered Species Act. N.S. Royal Gazette, Part II. Vol. 24, No. 13.
- PAPPAS, M.J. AND BRECKE, B.J. 1992. Habitat selection of juvenile Blanding's turtles, *Emydoidea blandingii*. *J. Herpetol.* 26:233-234.
- PETOKAS, P.J. 1986. Patterns of reproduction and growth in the freshwater turtle *Emydoidea blandingii*. Ph.D. Thesis, State Univ. New York, Binghamton.
- POWER, T.D. 1989. Seasonal movements and nesting ecology of a relict population of Blanding's turtle (*Emydoidea blandingii* (Holbrook)) in Nova Scotia. M.S. Thesis, Acadia University, Wolfville, Nova Scotia.
- POWER, T.D., HERMAN, T.B., AND KEREKES, J. 1994. Water colour as a predictor of local distribution of Blanding's turtles, *Emydoidea blandingii*, in Nova Scotia. *Can. Field-Nat.* 108:17-21.
- ROSS, D.A. 1989. Population ecology of painted and Blanding's turtles (*Chrysemys picta* and *Emydoidea blandingii*) in central Wisconsin. *Wisc. Acad. Sci. Arts Lett.* 77:77-84.
- ROSS, D.A. AND ANDERSON, R.K. 1990. Habitat use, movement, and nesting of *Emydoidea blandingii* in central Wisconsin. *J. Herpetol.* 24:6-12.
- SAJWAJ, T.D., PIEPGRAS, S.A., AND LANG, J.W. 1998. Blanding's turtle (*Emydoidea blandingii*) at Camp Ripley: critical habitats, population status, management guidelines. Final Report to Nongame Wildlife Office, Minnesota DNR, Brainerd, 179 pp.
- SLOAN, K.N., BUHLMANN, K.A., AND LOVICH, J.E. 1996. Stomach contents of commercially harvested adult alligator snapping turtles, *Macrolemys temminckii*. *Chelonian Conservation and Biology* 2:96-99.
- STANDING, K.L. 1997. Reproduction, nest site selection, and neonatal behaviour in a northern peripheral population of Blanding's turtle (*Emydoidea blandingii*). M.S. Thesis, Acadia University, Wolfville, Nova Scotia.
- STANDING, K.L., HERMAN, T.B., HURLBURT, D.D., AND MORRISON, I.P. 1997. Postemergence behaviour of neonates in a northern peripheral population of Blanding's turtle, *Emydoidea blandingii*, in Nova Scotia. *Can. J. Zool.* 75:1387-1395.
- TAYLOR, J. 1997. The status of turtles in New Hampshire. In: Tying, T.F. (Ed.). Status and Conservation of Turtles of the Northeastern United States. Lanesboro, MN: Serpent's Tale, pp. 4-5.

Received: 16 July 1997

Reviewed: 20 April 1998

Revised and Accepted: 25 May 1998